FAX FORM TO: 1.866.233.8317

PHONE: 1.855.492.0817

EMAIL: <u>contact@bluegrass-rx.com</u>

Complete the following <u>or include demographic sheet</u>.

1. PATIENT INFORMATION		2. PRESCRIBER INFORMATION					
Name:	Name:						
Address:		DEA #:	NPI #:	State Lic. #:			
City, State, ZIP:		Group or Hospital:					
Primary Phone: DOB: / /		Address:					
Alternate Phone: Gender:		City, State, Zip:					
Email:	Phone:	Fax:					
Primary Language: Last Fou	r of SSN:	Contact Person:	Phone:				
3. INSURANCE INFORMATION Fax copy of prescription and insurance cards with this form, if available (front and back)							
Primary Insurance Company Name:		Secondary Insurance Company Name:					
Primary Cardholder Name:		Secondary Cardholder Name:					
Relationship: 🔘 Self 🛛 Spouse/Partner 💭 Child/Dependent		Relationship: 🔵 Sel	Spouse/Partner	Child/Dependent			
Phone: Member ID:	Group #:	Phone:	Member ID:	Group #:			
4. DIAGNOSIS AND CLINICAL INFORMATION							
Needs by Date: / / Ship to: OPatient Office Other:							
Date of Diagnosis: / /	Type of therapy: O New O Continuing O Restart						
D59.5 Paroxysmal nocturnal hemoglobinuria (PNH)	Has patient received meningococcal vaccination? O Yes O No If no, reason:						
 D59.3 Hemolytic-uremic syndrome Other: 	Infusion appointment day: / / Time:						
	Allergies:						
Height (in/cm): Weight (lb/kg):	Current Medications:						

Medication	Dose/Strength	Directions	Quantity	Refills
Soliris 300 mg/30 ml vial (10 mg/ml)	For treatment of PNH:	4-week supply		
	O Dose titration - Month 1: Administer 600 mg via IV infusion every 7 days for 4 weeks	12-week supply		
	 Dose titration to maintenance - Month 2: Administer 900 mg via IV infusion every 2 weeks starting on week 5 			
	O Maintenance dosing: Administer 900 mg vial IV infusion every 2 weeks			
	O Other:			
		O For treatment of aHUS - 18 years or older	4-week supply	
		O Dose titration - Month 1: Administer 900 mg via IV infusion every 7 days for 4 weeks	12-week supply	
		 Dose titration to maintenance - Month 2: Administer 1200 mg via IV infusion every 2 weeks starting on week 5 		
		O Maintenance dosing: Administer 1200 mg vial IV infusion every 2 weeks		
		Other:		

Ancillary supplies and kits will be provided as needed for administration.

6. PRESCRIBER SIGNATURE								
x	/ /	x	/ /					
DISPENSE AS WRITTEN	DATE	PRODUCT SUBSTITUTION PERMITTED	DATE					

IMPORTANT NOTICE: This facsimile transmission is intended to be delivered only to the named addressee and may contain material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If it is received by anyone other than the named addressee, the recipient should immediately notify the sender at the address or telephone number set forth herein and obtain instructions as to disposal of the transmitted material. In no event should such material be read or retained by anyone other than the named addressee, except by express authority of the sender to the named addressee.